



Leukemoid Reaction in a Preterm Infant: Diagnostic Challenge in Resource Poor Setting: A Case Report

Okoli Chukwudi^{1*}, Nwaogu Nwaoma² and Nwafia Ifeyinwa³

¹Department of Haematology/Oncology, Lifeline Children's Hospital Lekki, Lagos, Nigeria.

²George's Memorial Medical Center Lekki, Lagos, Nigeria.

³College of Medicine, University of Nigeria Ituku, Ozalla Campus, Enugu, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Author OC designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors NN and NI managed the analyses of the study. Author OC managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJMPCR/2020/v13i230117

Editor(s):

(1) Dr. Jurij Janež, University Medical Centre Ljubljana, Slovenia.

Reviewers:

(1) Mariana Almada Bassani, University of Campinas, Brasil.

(2) Andreza de Lemos Bezerra, UNINASSAU, Brazil.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/61804>

Case Study

Received 25 July 2020

Accepted 30 September 2020

Published 19 October 2020

ABSTRACT

A preterm neonate delivered at 28 weeks gestation, mother had antenatal steroid. Blood counts showed leukemoid reaction, blood culture, procalcitonin and peripheral blood film was normal. Baby was stabilized in the NICU, recovered and the WBC count done serially showed a downward trend. The leukemoid reaction was presumed to come from antenatal steroid use. The diagnostic and management challenges encountered in managing the infant in resource constrained environment like ours is presented alongside.

Keywords: Leukemoid reaction; preterm infant; blood culture.

ABBREVIATIONS

NICU : Neonatal Intensive Care Unit;

WBC : White Cell Count;

CRP : C-Reactive Protein;

BMA : Bone Marrow Aspirate

*Corresponding author: E-mail: drchuksokoli@yahoo.com, chukwudi.okoli@npmcn.edu.ng;

1. INTRODUCTION

Leukemoid reaction is an extreme form of leukocytosis similar to that seen in leukemia but caused by other conditions. Leukocytosis exceeding 50,000 wbc/mm² with increase in early neutrophil precursors in the neonatal period is known as neonatal leukemoid reaction [1].

Leukemoid reaction in the neonatal period can be associated with sepsis, congenital leukemia, bronchopulmonary dysplasia, prematurity, antenatal steroid use and congenital abnormalities [2]. The incidence of leukemoid reaction in neonate range between 1.3% – 15% [3].

We report a case of leukemoid reaction in a preterm infant and the challenges in managing such diagnosis in resource poor environment.

2. CASE PRESENTATION

A preterm female baby was delivered caesarean section at 28 weeks + 4days on account of antepartum hemorrhage and preterm pre labour rupture of membranes. Apgar scores were 8 and 10 in the first and fifth minute respectively. Birth Weight was 1450 grams. Baby had respiratory distress and was admitted and nursed in an incubator with CPAP in the Neonatal Intensive care Unit. Investigations such as complete blood count (CBC), C-reactive protein, blood culture, procalcitonin, and peripheral blood film were sent. Intravenous antibiotic ceftazidime and amikacin was commenced and later stopped when blood culture did not reveal any organism. Babies initial investigation results; WBC 81,000 m/mm³, Neutrophils 66%, Lymphocytes 25%, monocytes 9% Hb 12.8g/dl, Platelets 311 m/mm³ on first day of life and by the 3rd day of life it was ; WBC 93,000 m/mm³, Neutrophils 74%, Lymphocytes 22.9%, monocytes 3.1% Hb 11.9g/dl, Platelets 260,000 m/mm³. The blood culture did not yield any organism, procalcitonine and CRP were normal while peripheral blood film did not show any abnormal cells and Lumber puncture was normal. Repeated CBC done alternate day showed a decreasing trend and by the 10th day post admission the blood count had reduced to, WBC 14,000 m/mm³, Neutrophils 33.83%, Lymphocytes 51.2%, monocytes 9.7% Hb 11.2 g/dl, Platelets 185,000 m/mm³ and remained within normal range till discharge. Bone marrow

aspiration and karyotype was not done. Baby continued to improve and was discharge home and has been seen for follow up with the complete blood count now normalized.

3. DISCUSSION

Leukocytosis is a common finding in newborns in the first few days of life [4]. This increase in leucocytes is from a surge or burst in cytokines (Granulocyte colony stimulating factor and Granulocyte-macrophage colony-stimulating factor) [5].

Leukemoid reaction have been demonstrated in up to 15% of preterm infants in the absence of any identifiable factor [5]. However it has been found to be more common in preterm infants, infections, antenatal steroid use, congenital leukemia and transient leukemoid reaction seen in Down syndrome [6].

In our reported case, patient's mother was on progesterone from the 7th – 15th week of pregnancy and had one dose of dexamethasone 48 hours prior to delivery, this history led credence to steroid use by mother being the cause of the leukemoid reaction in our case.

Our sepsis work-up in the patient was extensive despite the diagnostic difficulties and challenges faced especially in getting investigation results in real time in our environment.

Other diagnosis such as congenital leukemia was considered, however the peripheral blood film result did not reveal any abnormal cells or blast and the lactate dehydrogenase levels was not elevated. Bone marrow aspiration was not done due to financial challenges as patient could not afford it.

Our patient did not get the benefit of a karyotype as this investigation is not readily available in our environment and when available the turnaround time is in excess of 4 weeks. The index case did not have any dysmorphic features hence the possibility of transient myeloproliferative disease also known as transient abnormal myelopoiesis a form of leukemia seen in Down syndrome was excluded [7].

Leukemoid reaction is occurs in a different clinical scenario which includes prematurity, antenatal steroid use, infectious etiologies,

adverse reactions to medications, asplenia, congenital leukemia diabetic ketoacidosis, and septic shock with end-organ damage. Infants with trisomy 21 can often be born with a leukemoid reaction [8].

Evaluating and investigating leukemoid reaction involved elimination of the possible etiologies, by birth history to determine the gestational age and antenatal use, carrying out investigations which includes complete blood count, peripheral blood, C - reactive protein and blood culture to eliminate possibility of sepsis. Bone marrow aspirate to eliminate diagnosis of possible congenital leukemia and karyotyping for trisomy 21 [9,10].

The treatment of leukemoid reaction depends on the possible aetiology, more often than not the white cell counts would normalize as the aetiology treated [11,12].

The mechanism of steroid induced leukemoid reaction involves the following pathophysiology; steroids inhibit L-selectin synthesis at gene level, leading to demargination of leukocytes into the circulation, it also causes delayed migration of PMNs into tissues and rate of apoptosis, and release of non-segmented (band) neutrophils from bone marrow [13].

4. CONCLUSION

Diagnosing leukemoid reaction in preterm infants in very challenging due to lack of adequate diagnostic equipment, cost and prolonged turnaround time of the investigations. These limitations notwithstanding any WBC > 30,000 in any neonate should be thoroughly investigated to exclude the possibilities of sepsis, congenital leukemia and transient myeloproliferative disorders.

CONSENT

As per international standard or university standard, patients' written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Hill JM, Duncan CN. Leukemoid reactions. *Am J Med Sci.* 1941;201:847-857.
2. Leukemoid Holland P, Mauer AM. Myeloid leukemoid reactions in children. *Arch Pediatr Adolesc Med.* 1963;2013:568-75.
3. Blanchette V, Dror Y, Chan A. Hematology. in: MacDonald MG, Mullett MD, Seshia MM, editors. *Avery's neonatology – pathophysiology and management of the newborn.* 6th ed. Philadelphia: Lippincott Williams and Wilkins. 2005;1169–234.
4. Pesce MA. Reference ranges for laboratory tests and procedures. In: Kliegman RM, Behrman RE, Jenson HB, Stanton BF, editors. *Nelson Textbook of Pediatrics.* Philadelphia: Saunders. 2008; 2943–53.
5. Rastogi S, Rastogi D, Sundaram R, Kulpa J, Parekh AJ. Leukemoid reaction in extremely low birth weight infants. *Am J Perinatol.* 1999;16:93–7.
6. Sushanth, Avabrattha KS, Tauro KJ, Shwethadri GK. Hyperleukocytosis in a neonate: A diagnostic dilemma. *Indian J Med Paediatr Oncol.* 2010;31(3):86-88. DOI:10.4103/0971-5851.73596
7. Apollonsky N, Shende A, Ouansafi I, Brody J, Atlas M, Aygun B. Transient myeloproliferative disorder in neonates with and without Down syndrome: A tale of two syndromes. *J Pediatr Hematol Oncol.* 2008;30:860–4
8. Alatassi EU, Sukkar M, Garrada FN. Sepsis-induced Hyperleukocytosis in a Preterm. *Cureus.* 2019;11(9):e5594. DOI: 10.7759/cureus.5594
9. Jansen E, Emmen J, Mohns T, Donker, A. Extreme hyperleucocytosis of the premature. *BMJ Case Reports.* 2013; bcr2012008385. Available: <https://doi.org/10.1136/bcr-2012-008385>
10. Hoofien A, Yarden-Bilavski H, Ashkenazi S. *et al.* Leukemoid reaction in the pediatric population: etiologies, outcome, and implications. *Eur J Pediatr.* 2018; 177:1029–1036. Available: <https://doi.org/10.1007/s00431-018-3155-5>

11. Jain R, Bansal D, Marwaha RK. Hyperleukocytosis: Emergency management. Indian J Pediatr. 2013; 80: 144–8.
DOI:10.1007/s12098-012-0917-3
12. Yang L, Liu ZJ, Chen LQ, Hu JL, Zou N. Neonatal leukemoid reaction caused by enterococcus fecalis septicemia. Archives of Clinical and Medical Case Reports 4. 2020;086-090.
13. Steroid Induced Leukocytosis [Internet]. Epomedicine; 2016. [Cited 2020 Sep 26].

© 2020 Okoli et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/61804>